

## Isolation and Some Characterization of Vitellogenin and Its Related Egg Yolk Proteins from Coho Salmon (*Oncorhynchus kisutch*)

AKIHIKO HARA<sup>1</sup>, CRAIG V. SULLIVAN<sup>2</sup> and WALTON W. DICKHOFF<sup>3,4</sup>

<sup>1</sup>Nanae Fish Culture Experimental Station, Faculty of Fisheries, Hokkaido University, Nanae, Kameda, Hokkaido 041-11, Japan, <sup>2</sup>Department of Zoology, North Carolina State University, Raleigh, North Carolina 27695-7617, <sup>3</sup>School of Fisheries, University of Washington, Seattle, Washington 98195, <sup>4</sup>National Marine Fisheries Service, Northwest Fisheries Center 2725 Montlake Blvd. East, Seattle, Washington 98112, USA

**ABSTRACT**—Vitellogenin (Vg) and its related egg protein 1 (E1) and egg protein 2 (E2) were isolated from serum or eggs of mature female coho salmon by precipitation in distilled water followed by chromatography on Sepharose 6B (Vg) or Sephadex G-200 (yolk proteins). The coho salmon proteins reacted specifically with respective antisera raised against Vg (a-Vg), E1 (a-E1) or E2 (a-E2) purified from chum salmon. Two female-specific proteins were identified in serum from mature coho salmon. Coho salmon Vg had an apparent molecular weight of 540 kDa after chromatography on Superose 6, appeared as a major 240 kDa band and a minor 165 kDa band in SDS-PAGE, and resolved into a major 165 kDa band and several minor 70–150 kDa bands after SDS-PAGE under reducing conditions. It reacted immunologically with a-E1 and a-E2. The other female-specific serum protein, designated coho salmon E2, reacted with a-E2 but not with a-E1. The apparent molecular weight of purified E1 and E2 were 230 and 35 kDa, respectively, after chromatography on Superose 6. E1 appeared as two main bands of 150 and 92 kDa in SDS-PAGE which resolved into two smaller bands (92 kDa and 20 kDa) after reduction. E2 appeared as a 30 kDa band in SDS-PAGE and as a 15 kDa band after reduction. The above immunological and biochemical characteristics and subunit structure of coho salmon Vg, E1, and E2 were found to be nearly identical to the corresponding proteins in several other salmonid species of diverse genera. These properties of Vg have been highly conserved during salmonid evolution.

### INTRODUCTION

Vitellogenin (Vg) has been well-characterized in avian and amphibian species as a precursor for egg yolk. Hepatic synthesis of Vg is induced by estrogen in maturing females. The protein is released into the bloodstream from where it is taken up by developing oocytes and chemically modified in the process of yolk formation (for review, [24, 34, 36, 38]). For example, in the amphibian, *Xenopus*, Vg is a lipoglycophospho-protein complex of precursors to several individual egg yolk proteins including lipovitellin, phosvitin

and phosvettes [39].

In teleost fish, there are many reports of the identification of Vg or Vg-like proteins using immunological, electrophoretical and chromatographical methods [1, 3–5, 7–12, 25, 26, 32, 33, 35]. However, there are few reports on the specific biochemical relationship between serum Vg and the egg yolk proteins derived from it. We reported previously that salmonid Vg (female-specific serum protein) from rainbow trout, *Oncorhynchus mykiss* [16], chum salmon, *O. keta* [15] and white-spotted char, *Salvelinus leucomaenis* [17], is a precursor to at least two egg yolk proteins, egg protein one (E1) and egg protein two (E2). E1 was identified as a lipovitellin similar to that of *Xenopus* based on its physicochemical properties such as molecular weight, amino acid composition, sub-

Accepted November 13, 1992

Received September 7, 1992

<sup>1</sup> To whom all correspondence should be addressed.

unit structure and lipid content. Though E2 elutes in size exclusion chromatography in the fraction (mol. wt. 25–30 kDa) where Markert and Vanstone [22] reported two soluble egg yolk proteins that they termed phosvitin and the  $\beta'$ -component, we could identify only one antigenic protein in this fraction (E2). In addition, we identified two distinct female-specific proteins in serum from mature females or estrogen-treated males and immature fish using immunoelectrophoresis with rabbit antiserum raised against female-specific serum proteins from the species mentioned above [15, 17]. One of the mature female-specific serum proteins was identified as a Vg containing both E1 and E2 antigenicity. The other appeared to be a Vg fragment, since it was deficient in E1 antigenicity but had E2 antigenicity. We concluded that the two female-specific proteins found in the serum of maturing female salmonids represent a complex of E1 + E2 (Vg) and a free E2.

The present paper describes the identification of coho salmon (*Oncorhynchus kisutch*) Vg by immunological methods using antisera raised in rabbits against chum salmon Vg, E1, and E2, and the isolation and partial biochemical characterization of coho salmon Vg and its related egg yolk proteins.

## MATERIALS AND METHODS

### *Experimental animals, blood and tissue samples*

The experimental animals used in this study were adult female coho salmon that returned on their spawning migration to the University of Washington experimental fish hatchery where they were sampled for blood and eggs. They were fully mature but had not yet ovulated. Blood samples were collected from the caudal blood vessels using a syringe fitted with a 21 ga needle. The blood samples were held on ice and immediately transported to the laboratory where they were allowed to clot at 0–4°C overnight. The serum was collected after the blood samples were centrifuged at 1000×g for 15 min, and it was stored at –20°C until use.

One year-old immature coho salmon of both sexes were used for experiments in which Vg was

induced by estrogen treatment. They were anesthetized in a solution (50 mg/l) of tricaine methane sulfonate (Argent Chemical Lab.) buffered to pH 7.0 with sodium bicarbonate and injected intramuscularly with 1 mg of estradiol-17 $\beta$  per kg of body weight dissolved in 100% propylene glycol. Two weeks after hormone treatment, blood samples were collected from these fish and processed as described above. Ovulated eggs were collected from female coho salmon during their spawning season (November). The eggs were washed twice with 0.9% NaCl and then kept frozen at –20°C until use.

### *Purification of Vg and egg yolk proteins*

Coho salmon Vg and its related egg yolk proteins were isolated using our previously reported procedures [16]. Briefly, 2.5 ml of a sample of serum pooled from several mature females was added to 25 ml of ice-cold distilled-deionized water and the mixture was allowed to stand for 30 min at 0°C. A precipitate formed and was centrifuged at 2,500×g for 15 min at 4°C. The pellet was resuspended in water (25 ml), recentrifuged and dissolved in 0.5 ml of 0.02 M Tris-HCl buffer, pH 8.0, containing 2% NaCl and 0.1% NaN<sub>3</sub>. The solution was then applied to a 1.6×60 cm gel filtration column of Sepharose 6B (Pharmacia Inc.) equilibrated with the Tris-HCl buffer. The column flow rate was adjusted to 15 ml/hr and 1.9 ml fractions were collected. The protein concentration in the fractions was estimated by absorbance at 280 nm and the one major peak was collected as the purified Vg. The presence of Vg in the fractions was assessed by single radial immunodiffusion [21] using an antiserum raised in rabbits against chum salmon E1 [15].

Egg yolk was collected from ovulated eggs using a syringe fitted with a 22 ga needle. The yolk was centrifuged at 1000×g for 5 min and the supernatant was collected and added dropwise to a 10× volume of ice-cold distilled-deionized water. The precipitate that formed was sedimented by centrifugation at 1000×g for 30 min and the pellet was redissolved in Tris-HCl buffer. This procedure was repeated twice. The final clear solution in Tris-HCl buffer was applied to a 1.6×60 cm gel filtration column of Sephadex G-200 (Pharmacia

Inc.). The elution buffer, flow rate, fraction volume and assay of protein in the fractions during Sephadex G-200 chromatography were the same as described above for Sepharose 6B chromatography. The elution pattern yielded two peaks, designated as E1 and E2 according to our previously defined criteria [15, 17].

#### Antisera

Rabbit antisera raised against mature female chum salmon serum proteins, Vg and egg yolk proteins (E1 and E2) were prepared as described previously [15]. An antiserum was also raised in rabbits against the coho salmon egg yolk protein, E1, by intradermal injection with purified E1 (approximately 0.5 mg) emulsified in an equal volume of complete Freund's adjuvant. Injections were done 4 times at 7–10 day intervals.

#### Electrophoresis and immunological procedures

Immuno-electrophoresis and single radial immunodiffusion [21] and double immunodiffusion [28] were performed using 1.2% agarose gels as described previously [15]. The concentration of serum Vg was determined by single radial immunodiffusion using the antiserum to coho salmon E1 and the purified coho salmon Vg as the reference standard protein. Discontinuous polyacrylamide gradient (3.75–18.75%) gel electrophoresis (PAGE) was carried out with (SDS-PAGE) or without (Disc PAGE) addition of SDS to the gel and samples as described by O'Farrell [27]. Preparation of samples for SDS-PAGE was done as described previously [16]. The marker proteins used for molecular weight determination were lactate dehydrogenase (subunit, 36 kDa), egg albumin (45 kDa), catalase (subunit, 60 kDa), bovine serum albumin (67 kDa), ferritin (subunits, 18.5 and 220 kDa) and thyroglobulin (subunit, 330 kDa).

#### FPLC chromatography

Chromatography was performed on a Pharmacia FPLC system using a prepacked Superose 6 HR 10/30 column (Pharmacia Inc.). The column was used for determination of the molecular weights of the purified proteins and for observation of the elution pattern of serum proteins. Serum samples

were diluted 10× in Tris-HCl buffer and 200 µl of the solution, or about 6 mg of purified protein in 200 µl buffer, was applied to the column and eluted with the same buffer with a flow rate of 0.5 ml/min. The column was calibrated with the marker proteins: trypsinogen (24 kDa), egg albumin (45 kDa), bovine serum albumin (67 kDa), α-amylase (200 kDa), apoferritin (443 kDa) and thyroglobulin (669 kDa).

#### Comparative studies on salmonid Vg, E1 and E2

Blood serum and eggs were collected as described above from mature females of 10 species of salmonids from the genera *Salmo*, *Oncorhynchus*, *Salvelinus* (*Sl.*), and *Hucho*, which were obtained during annual spawning operations at various national, prefectural, university and commercial fish hatcheries in Hokkaido, Japan. These species included sockeye salmon (*O. nerka*), masu salmon (*O. masou*), amago salmon (*O. rhodurus*), rainbow trout, brown trout (*S. trutta*), Japanese huchen (*H. perryi*), brook trout (*Sl. fontinalis*), dolly varden (*Sl. malma*), white-spotted char (*Sl. leucomaenis*), and lake trout (*Sl. namaycush*). The egg yolk proteins, E1 and E2, were isolated as described above from the eggs of each of these species. The mature female serum, E1 and E2 of each species was reacted against the a-Vg, a-E1 and a-E2 antisera generated against the chum salmon proteins, and two additional a-Vg antisera generated against rainbow trout or spotted char vitellogenin [15, 16], in double immunodiffusion assays and immuno-electrophoresis as described above.

## RESULTS

#### Identification of coho salmon Vg

Immunological methods were used for identification of coho salmon Vg. Figure 1 shows immuno-electrophoresis of serum from vitellogenic female coho salmon using the four antisera against chum salmon proteins as follows: antiserum against mature female chum salmon serum proteins (a-S), specific antiserum to chum salmon Vg (a-Vg), and specific antisera to chum salmon egg yolk proteins E1 (a-E1) and E2 (a-E2). With the




FIG. 1. Immunoelectrophoresis of mature female coho salmon serum using four different antisera against chum salmon. a-S: antiserum to mature female chum salmon serum proteins, a-Vg: antiserum to chum salmon vitellogenin, a-E1: antiserum to egg yolk protein one (lipovitellin) of chum salmon, a-E2: antiserum to egg yolk protein two (phosvitin/ $\beta'$ -component fraction) of chum salmon.

exception of a-S, all the antisera reacted only with serum from vitellogenic females but not with male serum (data not shown). The a-Vg antiserum produced two distinct precipitin lines and the two antisera against egg yolk proteins (a-E1 and a-E2) each produced one precipitin line when reacted against serum from mature female coho salmon in immunoelectrophoresis. Based on our previous studies of salmonid Vg from rainbow trout and chum salmon [13] and white-spotted char [17], the two precipitin lines formed against mature coho salmon female serum by the a-Vg antiserum were considered to be Vg (the inner line formed near antigen well) and a fragment of Vg (the outer line formed near the antiserum trough), respectively. The Vg concentration in serum from mature but unovulated female coho salmon, assayed by single radial immunodiffusion, ranged between 4.68 mg/ml and 15.91 mg/ml (average 13.0 mg/ml).

#### *Purification of Vg and its related egg yolk proteins*

The elution pattern in Sepharose 6B chromatography of protein from the water-insoluble fraction of coho salmon serum is shown in Fig. 2. The main peak was found to consist of Vg when assayed by single radial immunodiffusion using the a-E1 antiserum. A single and symmetrical peak

was obtained and collected as the purified coho salmon Vg. After immunoelectrophoresis, this preparation of Vg gave rise to only one precipitin line when reacted against the a-S and a-Vg antisera, as shown in Fig. 3. As shown in Fig. 4, the elution pattern of the water-insoluble fraction of coho salmon egg extracts consisted of two distinct peaks, designated E1 and E2, after gel filtration on Sephadex G-200. The relative protein concentrations of E1 and E2, determined planimetrically from charts of gel filtration monitored by absorption at 280 nm, were found to be 13.3 and 1.0 respectively. When adjusted for the estimated molecular weights of E1 and E2 (discussed below), the apparent molar ratio of E1 to E2 is 2 to 1. The peak fractions were tested for their reactivity to the a-Vg, a-E1 and a-E2 antisera by immunodiffusion. The fraction designated coho salmon E1 reacted with the a-Vg antisera and a-E1 antisera, but not with antiserum a-E2. The fraction designated coho salmon E2 reacted with the a-Vg and a-E2 antisera but not with antiserum a-E1 (data not shown). After Disc-PAGE, the preparation of coho salmon Vg displayed a major band and a secondary band which migrated faster than the major band as shown in Fig. 5 (lane e). Some faint bands that migrated more slowly than the major




FIG. 2. Elution pattern of the water-insoluble fraction of coho salmon serum chromatographed on Sepharose 6B. The fractions pooled from tube number 43-48 were collected and concentrated as coho salmon vitellogenin for further analysis.



FIG. 3. Immunoelectrophoresis of purified coho salmon vitellogenin. S: mature female coho salmon serum, Vg: purified coho salmon vitellogenin, a-Vg and a-S: same as in Fig. 1.




FIG. 4. Chromatography on Sephadex G-200 of water-insoluble proteins isolated from coho salmon eggs. Two major peaks were designated as E1 and E2.

band were also observed. The major band of purified Vg migrated to the position corresponding to the strong band observed only in serum of mature females (lane b) and estradiol-treated fish (lane d). While coho salmon E1 gave a single distinct band in Disc-PAGE (Fig. 5, lane g), coho salmon E2 gave a rather broad band composed of several sharp lines (Fig. 5, lane h). These results are essentially identical to those we reported for chum salmon [15].

In order to compare antigenicity between the three purified coho salmon proteins, double immunodiffusion was performed in 1.2% agarose gels using the  $\alpha$ -Vg antiserum. The precipitin line from coho salmon Vg formed a spur over the precipitin lines of the two egg yolk proteins, E1 and E2, and the precipitin lines of each egg yolk

protein crossed each other (data not shown), confirming that the coho salmon Vg molecule contains both E1 and E2 antigenicity.

#### *Molecular weight of purified proteins*

As shown in Fig. 6, the molecular weights of the purified coho salmon proteins, estimated by FPLC chromatography on the Superose 6 column, were 540 kDa for Vg, 230 kDa for E1, and 35 kDa for E2. The electrophoretic patterns of coho salmon Vg, E1 and E2 after SDS-PAGE  $\pm$  reduction with 2-mercaptoethanol (2-ME) are shown in Fig. 7. Vg displayed one main band and a minor band corresponding to molecular weights of 240 kDa and 165 kDa, respectively (lane C). Vg reduced with 2-ME showed a main band with a molecular weight of 165 kDa as well as some minor bands of lower




Fig. 5. Gradient Disc-PAGE of purified vitellogenin, E1 and E2 and egg yolk extracts from coho salmon. a: serum proteins from mature male, b: serum proteins from mature female, c: serum from immature male (control fish from estrogen-treatment experiment), d: serum from estrogen-treated immature fish, e: vitellogenin, f: egg yolk extracts, g: E1, h: E2.

(70 to 150 kDa) molecular weight (lane c). E1 showed a main band corresponding to a molecular weight of about 150 kDa and minor 92 kDa band, as well as a faint band migrating near the dye front (lane D). After reduction with 2-ME, E1 appeared as two main 92 kDa and 20 kDa bands, as well as some faint bands of intermediate (70 kDa to 90 kDa) molecular weight. E2 appeared in SDS-PAGE as a single 30 kDa band before reduction, and as a single 15 kDa band after reduction with 2-ME. These results were very similar to those observed Vg, E1 and E2 of rainbow trout and chum salmon [15].

#### *Serum elution profiles in FPLC*

Typical patterns of coho salmon serum after

chromatography on Superose 6 are shown in Fig. 8. Distinct differences between the sexes were seen in the patterns. The major peak from female serum eluting at effluent volume 13.9 ml (panel b) was not seen in male serum (panel a). This peak was also induced in immature fish by treatment with estradiol-17 $\beta$  (panel c). Purified Vg (see Fig. 6) eluted the same position (arrow, Fig. 8) as the mature female specific and estradiol-inducible peak.

#### *Cross-reactivity of antisera to chum salmon with other salmonid fish*

The elution patterns on Sephadex G-200 of E1 and E2 from the 10 species of salmonids were nearly identical to that seen for coho salmon (data




FIG. 6. Estimation of the molecular weight of coho salmon vitellogenin by Supurose 6 gel filtration in the FPLC system. The open circles correspond to the elution volume of purified vitellogenin (Vg), E1 and E2. The marker proteins used for calibration and indicated by the dark circles were, 1: trypsinogen, 2: egg albumin, 3: bovine serum albumin, 4:  $\alpha$ -amylase, 5: apoferritin, and 6: thyroglobulin.




FIG. 7. SDS-PAGE on 3.75–18.75% gradient slab gels of mature male serum (A, a), female serum (B, b), vitellogenin (C, c), E1 (D, d), and E2 (E, e) in coho salmon. Lane A-E represent non-reduced samples and lanes a-e represent samples reduced with 2-mercaptoethanol.






FIG. 8. Elution profiles of mature male (a) and female serum (b), and estrogen-treated male serum (c) in coho salmon on Supurose 6 in the FPLC system. The arrow indicates the elution position of vitellogenin.

not shown). As noted above, the a-Vg, a-E1 and a-E2 antisera raised against chum salmon Vg, E1 and E2, respectively, reacted strongly and specifically to the respective coho salmon proteins. In addition, we found that these antisera, as well as two other a-Vg antisera raised against Vg from rainbow trout and white-spotted char [15, 16], yielded a pattern of immunoreactivity identical to that reported above for coho salmon when they were reacted in double immunodiffusion assays and immunoelectrophoresis against the mature female serum, E1 and E2, from the 10 salmonid species. For each species, two mature female-specific serum proteins were seen in immunoelectrophoresis. One reacted with all of the antisera tested and the other reacted only with the a-Vg and a-E2 antisera. E1's from the various species

reacted with all three a-Vg's and with a-E1, but not with a-E2. E2's from the various species reacted with all three a-Vg's and with a-E2, but not with a-E1.

## DISCUSSION

In the present study, coho salmon Vg was purified within one day from the serum of mature females by precipitation with distilled water followed by gel filtration on Sepharose 6B. When the gel filtration step was done on a Supurose 6 column using the FPLC system, complete purification of Vg was accomplished within 3 hours, but yields were much reduced (data not shown). The precipitation step with distilled water is a very simple, easy and effective method for purification of sal-

monid Vgs, though its success depends on the starting conditions of the serum and its concentration of Vg. For example, precipitation of coho salmon Vg was not possible using serum containing Vg concentrations below about 5 mg/ml or using serum that was not previously frozen. We have used hydroxylapatite column chromatography to purify Vg from salmonid serum with low Vg concentrations that cannot be precipitated with distilled water, and the purified protein is similar in all respects to that obtained using the precipitation method [14–17].

So far, the detection of teleost Vg has been performed primarily by immunological, electrophoretic and chromatographical procedures. In the present study, in addition to immunological methods, we identified Vg in serum of mature females and estrogen-treated males and immature fish using a Supurose 6 column in an FPLC chromatography system. This method required only one drop of serum for assay and the procedure is simple and rapid. Silversand and Haux [31] recently reported a similar method for purification of turbot Vg, using a Mono Q column in an FPLC system. Though further studies will be needed to confirm the relation between the concentration of Vg and its peak height or area in FPLC or HPLC chromatography, the method should prove useful for rapid routine assay of fish Vgs.

Vertebrate Vg has been generally confirmed to be the precursor of the major yolk proteins, lipovitellin and phosvitin. Rainbow trout egg yolk proteins can be chromatographically separated into three major components [20]. Phosvitin is a phosphoprotein rich in serine [5, 19, 30], soluble in trichloroacetic acid [5, 30, 37], and poorly stained with Coomassie Blue on electrophoretic gels [18]. Lipovitellin is the main yolk protein of oviparous vertebrates, and it contains a high concentration of lipids and little or no protein phosphorus. In rainbow trout, the lipovitellin has been reported to consist of two subunits of mol. wt. 90 kDa and 15 kDa [16] or 95 and 24 kDa [6] or 92 and 20 kDa [1], or to consist of four subunits from 145–160 kDa ( $n=2$ ) and from 14–19 kDa ( $n=2$ ) which can easily dimerize [29]. A third yolk protein, the  $\beta'$ -component [20], is characteristic of the yolk of salmonid eggs [22, 23]. Its amino acid composition

is distinct from those of lipovitellin and phosvitin [5]. It can escape precipitation when the yolk is diluted with distilled water [20, 22]. Though the origin of the  $\beta'$ -component remains to be conclusively verified, it has been suggested that it is derived from Vg like lipovitellin and phosvitin [38]. In the present study, the water-insoluble egg yolk proteins of coho salmon were separated into two components, namely, E1 and E2. The molecular weight of E1 and E2 were estimated by gel filtration to be 230 kDa and 35 kDa, respectively. Using SDS-PAGE without reduction by 2-ME, the molecular weights of E1 and E2 were estimated to be 150 kDa and 30 kDa, respectively. Upon reduction with 2-ME, the 150 kDa component of E1 further split into two major polypeptides with molecular weights of 92 and 20 kDa. The 30 kDa E2 split into a single component with a molecular weight of 15 kDa, suggesting that it may be a homodimer. These results are essentially identical to those that we previously reported for rainbow trout egg yolk proteins [16].

Immunological methods using antisera to chum salmon Vg, E1 and E2 revealed two female-specific proteins in the serum of mature coho salmon. One (Vg) was, in part, a complex of E1 + E2. The observation that coho salmon Vg contained both the E1 and E2 components of egg yolk proteins is similar to our previously reported results on other salmonid species [15]. The second female-specific serum protein contained only E2 antigenicity. It did not react with antiserum to E1. A relation of these two serum proteins to the coho salmon  $\beta'$ -component and/or phosvitin [23] remains to be established. Our results suggest that an antiserum raised against purified E1 would be preferable to an antiserum to Vg for measurement of Vg concentrations in serum.

Chen et al. [7] developed a rapid sensitive assay for rainbow trout Vg using an antiserum against lipovitellin in rocket immunoelectrophoresis. They reported that their antibody against rainbow trout lipovitellin showed a continuous precipitation arc when reacted against coho salmon lipovitellin in Ouchterlony immunodiffusion and Western immunoblotting assays. These results support our observation that antisera to chum salmon Vg, E1 and E2 strongly react to the

respective coho salmon proteins. We also found that the antiserum to chum salmon Vg, E1 and E2, react fully and specifically with the corresponding proteins of various other species from diverse salmonid genera, including sockeye salmon, masu salmon, amago salmon, rainbow trout, brown trout, Japanese huchen, brook trout, dolly varden, white-spotted char, and lake trout. Benfey et al. [2] also showed that plasma from various salmonid species exhibited parallelism in a coho salmon Vg radioimmunoassay. These results suggest that affinity chromatography on a column containing antibodies against the proteins purified from one salmonid species should be a viable approach to purifying Vg from a variety of other salmonid fishes. They also indicate that Vg and its related egg proteins have been highly conserved during salmonid evolution.

#### ACKNOWLEDGMENTS

We would express our gratitude to C. V. W. Mahnken, National Marine Fisheries Service, Northwest Fisheries Center, Seattle, Washington, for facilities and financial support, and to Drs. E. Brannon and W. K. Hershberger, School of Fisheries, University of Washington, Seattle, Washington, for access to fish and samples. Appreciation is extended to Drs. E. Plisetskaya and P. Swanson, School of Fisheries, University of Washington, for their kind advice and valuable suggestions, to M. G. Bernard and Dr. L. Yan for help with fish sampling and care, and to Dr. K. Takano for help collecting samples from the various species of salmonids. This investigation was supported by grant from the National Science Foundation (DCB 8615521) to W.W.Dickhoff, by a grant from the International Exchange Programs Committee of Hokkaido University, Sapporo, Japan, to C.V.Sullivan, and by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Education, Science and Culture (01440014) to A. Hara.

#### REFERENCES

- 1 Babin PJ (1987) Apolipoproteins and the association of egg yolk proteins with plasma high density lipoproteins after ovulation and follicular atresia in the rainbow trout (*Salmo gairdneri*). *J Biol Chem* 262: 4290-4296
- 2 Benfey TJ, Donaldson EM, Owen TG (1989) An homologous radioimmunoassay for coho salmon (*Oncorhynchus kisutch*) vitellogenin, with general applicability to other Pacific salmonids. *Gen Comp Endocrinol* 75: 78-82
- 3 Bradley JT, Grizzle JM (1989) Vitellogenin induction by estradiol in channel catfish, *Ictalurus punctatus*. *Gen Comp Endocrinol* 73: 28-39
- 4 Campbell CM, Idler DR (1976) Hormonal control of vitellogenesis hypophysectomized winter flounder (*Pseudopleuronectes americanus* Walbaum). *Gen Comp Endocrinol* 28: 143-150
- 5 Campbell CM, Idler DR (1980) Characterization of an estradiol-induced protein from rainbow trout as vitellogenin by the cross reactivity to ovarian yolk fractions. *Biol Reprod* 22: 605-617
- 6 Chen TT (1983) Identification and characterization of estrogen-responsive gene products in the liver of rainbow trout. *Can J Biochem Cell Biol* 61: 802-810
- 7 Chen TT, Reid PC, Van Beneden R, Sonstegard RA (1986) Effect of Aroclor 1254 and Mirex on estradiol-induced vitellogenin production in juvenile rainbow trout (*Salmo gairdneri*). *Can J Fish Aquat Sci* 43:169-173
- 8 Copeland PA, Sumpter JP, Walker TK, Croft M (1986) Vitellogenin levels in male and female rainbow trout (*Salmo gairdneri* Richardson) at various stages of the reproductive cycle. *Comp Biochem Physiol* 83B: 487-493
- 9 Covens M, Covens L, Ollevier F, de Loof A (1987) A comparative study of some properties of vitellogenin (Vg) and yolk proteins in a number of freshwater and marine teleosts. *Comp Biochem Physiol* 88B: 75-80
- 10 de Vlaming VL, Wiley HS, Delahunty G, Wallace RA (1980) Goldfish (*Carassius auratus*) vitellogenin: Induction, isolation, properties and relationship to yolk proteins. *Comp Biochem Physiol* 67B: 613-623
- 11 Emmersen BK, Petersen IM (1976) Natural occurrence and experimental induction by estradiol 17- $\beta$ , of a lipophosphoprotein in flounder (*Platichthys flesus* L.). *Comp Biochem Physiol* 54B: 443-446
- 12 Fletcher PE, Fletcher GL (1980) Zinc- and copper-binding proteins in the plasma of winter flounder (*Pseudopleuronectes americanus*). *Can J Zool* 58: 609-613
- 13 Hara A (1976) Iron-binding activity of female-specific serum proteins of rainbow trout (*Salmo gairdneri*) and chum salmon (*Oncorhynchus keta*). *Biochim Biophys Acta* 427: 549-557
- 14 Hara A (1978) Sexual differences in serum proteins of chum salmon and the purification of female-specific serum protein. *Bull Jap Soc Sci Fish* 44: 689-693
- 15 Hara A (1987) Studies on female-specific serum proteins (vitellogenin) and egg yolk proteins in teleosts:Immunochemical, physicochemical and structural studies. *Mem Fac Fish Hokkaido Univ* 34: 1-59
- 16 Hara A, Hirai H (1978) Comparative studies on

- immunochemical properties of female-specific serum protein and egg yolk proteins in rainbow trout (*Salmo gairdneri*). *Comp Biochem Physiol* 48B: 389-399
- 17 Hara A, Matsubara T, Saneyashi M, Takano K (1984) Vitellogenin and its derivatives in egg yolk proteins of white-spotted char (*Salvelinus leucomaenis*). *Bull Fac Fish Hokkaido Univ* 35: 144-153
  - 18 Hegenauer J, Ripley L, Nace G (1977) Staining acidic phosphoproteins (phosvitin) in electrophoretic gels. *Anal Biochem* 78: 308-311
  - 19 Ito Y, Fujii T, Kanamori M, Hattori T, Yoshioka R (1966) Phosvitin of the trout roe. *J Biol Chem* 60: 726-728
  - 20 Jared DW, Wallace RA (1968) Comparative chromatography of the yolk proteins of teleosts. *Comp Biochem Physiol* 24: 437-443
  - 21 Mancini G, Carbonara AO, Heremans JF (1965) Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochemistry* 2: 235-254
  - 22 Markert JR, Vanstone WE (1968) Egg proteins of Pacific salmon, genus *Oncorhynchus*: Proteins in the soluble fraction. *Can J Biochem* 46: 1334-1337
  - 23 Markert JR, Vanstone WE (1971) Egg proteins of coho salmon (*Oncorhynchus kisutch*): chromatographic separation and molecular weights of the major proteins in the high density fraction and their presence in salmon plasma. *J Fish Res Board Can* 28: 1853-1856
  - 24 Mommsen TP, Walsh PL (1988) Vitellogenesis and oocyte assembly. In "Fish Physiology Vol XIA" Eds by WS Hoar, DJ Randall, Academic Press, New York, pp 347-406
  - 25 Nath P, Sundararaj BI (1981) Isolation and identification of female-specific serum lipophosphoprotein (vitellogenin) in the catfish, *Heteropneustes fossilis*. *Gen Comp Endocrinol* 43: 184-190
  - 26 Ng TB, Idler DR (1983) Yolk formation and differentiation in teleost fishes. In "Fish Physiology Vol IXA" Eds by WS Hoar, DJ Randall, Academic Press, New York, pp 373-404
  - 27 O'Farrell PH (1975) High resolution two dimensional electrophoresis of proteins. *J Biol Chem* 250: 4007-4021
  - 28 Ouchterlony Ö (1953) Antibody-antigen reactions in gels. IV. Types of reactions in coordinated systems of diffusion. *Acta Path Microbiol Scand* 32: 231-240
  - 29 Riaz A, Fremont L, Gozzelino MT (1988) Characterization of egg yolk proteins from rainbow trout, *Salmo gairdneri* (Rich). *Comp Biochem Physiol* 89B: 399-407
  - 30 Schmidt G, Bartsch G, Kitagawa T, Fujisawa K, Knolle J, Joseph J, De Marco P, Liss M, Haschemeyer R (1965) Isolation of a protein of high phosphorus content from the eggs of brown brook trout. *Biochem Biophys Res Commun* 18: 60-65
  - 31 Silversand C, Haux C (1989) Isolation of turbot (*Scophthalmus maximus*) vitellogenin by high-performance anion-exchange chromatography. *J Chromatography* 478: 387-397
  - 32 So YP, Idler DR, Hwang SJ (1985) Plasma vitellogenin in landlocked Atlantic salmon (*Salmo salar* Quananche): Isolation, homologous radioimmunoassay and immunological cross-reactivity with vitellogenin from other teleosts. *Comp Biochem Physiol* 81B: 63-71
  - 33 Sullivan CV, Tao Y, Hodson, R.G, Hara A, Bennett RO, Woods III LC (1991) Vitellogenin and vitellogenesis in striped bass (*Morone saxatilis*) broodstock. In "Proceedings of the Fourth International Symposium on the Reproductive Physiology of Fish". Eds by AP Scott, JP Sumpter, DE Kime, MS Rolfe, Sheffield, U.K. pp 315-317
  - 34 Tyler C (1991) Vitellogenesis in salmonids. In "Proceedings of the Fourth International Symposium on the Reproductive Physiology of Fish". Eds by AP Scott, JP Sumpter, DE Kime, MS Rolfe, Sheffield, U.K. pp 295-299
  - 35 Wallace RA (1978) Oocyte growth in nonmammalian vertebrates. In "The Vertebrate Ovary". Ed by RE Jones, Plenum Press, New York, pp 469-502
  - 36 Wallace RA (1985) Vitellogenesis and oocyte growth in nonmammalian vertebrate. In "Development Biology Vol 1" Ed by LW Browder, Plenum Press, New York, pp 127-177
  - 37 Wallace RA, Jared DW, Eisen AZ (1966) A general method for the isolation and purification of phosvitin from vertebrate eggs. *Can J Biochem* 44: 1647-1655
  - 38 Wiegand MD (1982) Vitellogenesis in fishes. In "Reproductive Physiology of Fish". Eds by C.J.J. Richter, HJT Goos, Wageningen, pp 136-146
  - 39 Wiley HS, Wallace RA (1981) The structure of vitellogenin: Multiple vitellogenins in *Xenopus laevis* give rise to multiple forms of the yolk proteins. *J Biol Chem* 256: 8626-8634